

# EFFECTS OF DRYING TEMPERATURE ON PROXIMATE COMPOSITION AND FUNCTIONAL PROPERTIES OF (*Colocasiaesculenta*) COCOYAM FLOUR

OGUNDELEOLUSOLA D<sup>\*</sup>, THOMPSON SAMUEL O<sup>\*</sup>,  
LAWALSONSIMISOLA K<sup>\*</sup>, DEMEHIN BLESSING F<sup>\*</sup>

## ABSTRACT

This research evaluated the effects of drying temperature on the proximate composition and functional properties of (*Colocasiaesculenta*) cocoyam flour. Flour was obtained from *Colocasiaesculenta* by peeling, slicing, washing, drying at different temperature (50 °C, 60 °C, 70 °C, 80 °C, and 90 °C), milling and sieving. The flours obtained were evaluated for proximate composition and functional properties. The flour samples had proximate composition ranging from 5.98 % to 11.28 % moisture content, 0.75 % to 3.22 % ash content, 4.93 % to 7.26 % crude protein, 0.41 % to 0.93 % fat content, 2.75 % to 3.19 % crude fiber and 72.69 % to 81.23 % carbohydrate. The flour samples have functional properties ranging from 1.32 mL/g to 2.90 mL/g water absorption capacity, 0.63 mL/g to 1.43 mL/g oil absorption capacity, 0.59 g/mL to 0.95 g/mL bulk density, 81 °C to 90°C gelatinization point, 14.30 to 18.60swelling capacity, 12 secs to 23 secs wett ability and 21 cp to 35cp viscosity measurement. The result of this study showed that the drying temperature have effect on both the proximate composition and functional properties of (*Colocasiaesculenta*) cocoyam flour.

**KEYWORDS:** *Colocasiaesculenta*, Proximate Composition, Functional Properties, Temperature, Drying.

## INTRODUCTION

Food nutrients majorly include carbohydrate, lipids, proteins, minerals, vitamins and water which can be derived from plant and animals. Animals flesh are consumed as food while plant is the only living organism that is capable of carrying out photosynthesis which makes them grow and plants are the major source of food in nature's chain [1]. Food originated from plant may be classified into cereals, roots and tubers,

sugars and syrups, legumes, pulses, nuts and oil seeds, vegetables and fruits [2].

Roots and tubers used as food include yams, cassava, potatoes, sweet potatoes and cocoyam. Cocoyams are monocotyledonous herbs that belong to the family Araceae and are grown primarily for their roots which are edible.

---

\*Chemistry Department, Federal University of Technology, Akure, Nigeria.

Correspondence E-mail Id: editor@eurekajournals.com

The name cocoyam is generally applied to a variety of useful and edible species belonging to different genera including *Colocasia*, *Xanthosoma*, *Alocasia*, *Crytospema* and *Amorphophallus* [3]. They are the third most important root crop (after yam and cassava), one of the most valuable root crops cultivated in Nigeria, and the second most valuable in West Africa. By far, more important and more extensive cultivation in Nigeria are *Colocasia* and *Xanthosoma*. Nigeria has the largest population of cocoyam consumers, followed by Ghana. The Southern part of Nigeria is known for bulk production of cocoyam [4].

Roots and tubers contribute about 20-48% of the total calories and about 7.1% protein to the diets of the people of sub-Saharan Africa. In Nigeria, they are the main sources of calories accounting for over 50% of the caloric intake of the people of the south. Yam has less than 6% protein while cassava is a poor source of protein, less than 3%. Cocoyam is fair in protein, 7-9% and calcium while sweet potatoes are poor in protein [5].

However, considering the nutritional quality of cocoyam, the high starch content and its quality (*i.e.* fine starch grains), the level of utilization of

cocoyam and its products both domestically and industrially is quite low. Cocoyam has high moisture content resulting in short storage life under ambient conditions and because of these, they are highly perishable and huge losses can occur after harvest [6]. There is a need to investigate the effect of temperature on the properties of cocoyam in order not to denature and devalue the nutrient in cocoyam while processing for storage.

## **MATERIALS AND METHODS**

Fresh corm of cocoyam (*Colocasia esculenta*) was purchased from a local market in Ondo State. All reagent and solvent that were used are of analytical grade.

## **SAMPLE PREPARATION**

The fresh corms of cocoyam (*Colocasia esculenta*) was washed, peeled, and then rinsed under running water. It was then sliced into 5mm thickness and the pieces were dried in the oven at 50°C, 60°C, 70°C, 80°C, and 90°C for 3 hours. The samples were sieved and packaged in air tight properly labeled polythene sachets for further analysis.



**Figure 2.1. *Colocasia esculenta* leaves**



Figure 2.2. *Colocasia esculenta* tubers

### PROXIMATE ANALYSIS

The proximate analysis was determined using the method described by [8]

#### DETERMINATION OF MOISTURE CONTENT

Exactly 2 g each of the samples were weighed and dried at about 105 °C in the oven for four hours to constant weight. The moisture content was reported as percentage loss in weight [8].

$$\text{Moisture content} = \frac{W_2 - W_3}{W_2 - W_1} \times 100 \dots \text{Equation 1}$$

Where:  $W_1$  = Weight of empty dish.

$W_2$  = Weight of dish and sample before drying

$W_3$  = Weight of dish and sample after drying

#### DETERMINATION OF ASH CONTENT

The ash content was determined by igniting 5 g of dry sample in a muffle furnace at about 550 °C to constant weight. It was cooled in a desiccator and weighed. The ash content was reported as a percentage dry mass [9].

$$\text{Percentage Ash} = \frac{\text{Weight loss } (W_2 - W_3)}{\text{Weight of sample } (W_1)} \times 100 \quad \text{Equation 2}$$

Where:  $W_1$  = weight of sample

$W_2$  = weight of sample + crucible

$W_3$  = weight of sample + crucible (constant weight after drying)

#### DETERMINATION OF CRUDE PROTEIN

Exactly 5 g of the samples were weighed and digested in macro-Kjeldahl apparatus with concentrated sulphuric acid. The ammonia liberated from the resulting ammonium sulphate after adding sodium hydroxide was distilled into 1 M boric acid and then titrated with 0.1 M HCl. The nitrogen value estimated was multiplied by 6.25 (protein factor) to obtain the value of the crude protein, expressed as the percentage of sample mass [9].

$$\% \text{ Nitrogen} = \frac{\text{Vol. of acid used} \times \text{Molarity} \times 0.014 \times \text{Dilution factor}}{\text{Weight of sample}} \times 100 \quad \text{Equation 3}$$

$$\% \text{ Protein} = \% \text{ N} \times 6.25 \quad \text{Equation 4}$$

#### DETERMINATION OF FAT

The crude fat was extracted from 5 g of each sample using a solvent extraction apparatus (Soxhlet apparatus) with low boiling point petroleum ether as solvent. The weight of the lipid obtained after evaporating off the solvent

from the extract gave the weight of the lipid present in the sample [9].

$$\text{Percentage of Crude fat} = \frac{\text{Weight loss } (W_2 - W_3)}{\text{Weight of sample } (W_1)} \times 100 \quad \text{Equation 5}$$

Where:  $W_1$  = weight of sample

$W_2$  = weight of sample + filter paper

$W_3$  = weight of sample + filter paper (constant weight after drying)

#### **DETERMINATION OF CRUDE FIBRE**

Exactly 5 g of sample were weighed and exhaustive extraction of substances soluble in 1.25% boiling sulphuric acid and 1.25% boiling sodium hydroxide was employed. The residual matter of crude fibre and inorganic material recovered and ash yielded the crude fibre expressed as percentage loss in weight of ashed residue [9].

$$\text{Percentage of Crude fibre} = \frac{\text{Weight loss } (W_2 - W_3)}{\text{Weight of sample } (W_1)} \times 100 \quad \text{Equation 6}$$

Where:  $W_1$  = Initial weight of sample

$W_2$  = weight of sample + crucible before ashing

$W_3$  = weight of sample + crucible after ashing (constant weight after drying).

#### **CARBOHYDRATE CONTENT DETERMINATION**

Carbohydrate content was determined by difference. The percentage total carbohydrate is estimated to be equal to the sum of percentage moisture, protein, ash and fibre subtracted from 100.

$$\% \text{ Carbohydrate} = 100 - (\% \text{ protein} + \% \text{ fat} + \% \text{ fibre} + \% \text{ ash} + \% \text{ moisture}) \quad \text{Equation 7}$$

#### **DETERMINATION OF FUNCTIONAL PROPERTIES**

The method of [10] was used for the determination of functional properties. The functional properties determined include water

and oil absorption capacities (1g sample); bulk density (5g sample); swelling index (3g sample) and wett ability capacity (1g sample). Gelatinization temperature was determined using the method described by [10].

#### **WATER / OIL ABSORPTION CAPACITY**

Exactly 1 g of sample was weighed into a clean conical graduated centrifuge tube and was mixed thoroughly with 10 mL distilled water/oil using a warring mixer for 30 secs. The sample was then allowed to stand for 30 mins at room temperature, after which it was centrifuged at 1000 rpm for 30 mins. After centrifugation, the volume of the free water (supernatant) or oil was read directly from the graduated centrifuge tube. The absorbed water was converted to weight (in grams) by multiplying by the density of oil (0.894 g/mL) and water (1 g/mL). The oil and water absorption capacities were expressed in grams of oil/water absorbed per gram of flour sample.

$$\text{Absorbed water} = \text{total water} - \text{free water} \quad \text{Equation 8}$$

#### **BULK DENSITY**

The gravimetric method was used. A weighed sample (10 g) was put in a calibrated 10 mL measuring cylinder. Then the bottom of the cylinder was tapped repeatedly onto a firm pad on a laboratory bench until a constant volume was observed. The packed volume was recorded. The bulk density is calculated as the ratio of the sample weight to the volume occupied by the sample after tapping [11].

$$\text{Bulk density (g/mL)} = \frac{\text{Weight of Sample (g)}}{\text{Volume of sample (mL)}} \quad \text{Equation 9}$$

#### **GELATINIZATION POINT**

Exactly 10 g of flour sample was suspended in distilled water in a 250 mL beaker and made up to 100 mL flour suspension. The aqueous suspension was heated in a boiling water bath,

with continuous stirring using a magnetic stirrer. A thermometer was then clamped on a retort stand with its bulb submerged in the suspension. The heating and stirring continued until the suspension began to gel and corresponding temperature was recorded 30 secs after gelatinization was visually noticed.

### **SWELLING CAPACITY**

This was determined as the ratio of the swollen volume to the ordinary volume of a unit weight of the flour. The method of [11] was used. One gram (1 g) of the sample was weighed into a clean dry measuring cylinder. The volume occupied by the sample was recorded before 5 mL of distilled water was added to the sample. This was left to stand undisturbed for an hour, after which the volume was observed and recorded again. The capacity of swelling ability of the sample was given by the formula below:

$$\text{Swelling Capacity} = \frac{\text{Volume occupied by sample after swelling}}{\text{Volume occupied by sample before swelling}}$$

### **WETTABILITY**

The method described by [12] was used. A graduated cylinder (25 mL) was washed and dried in the oven and one gram of sample was weighed out and filled into the cylinder. A 600 mL beaker was filled with distilled water up to the 500 mL mark. A finger was placed over the open end of the 25 mL cylinder containing the sample. The cylinder was inverted and clamped at a height of 10 cm from the surface of a 600 mL beaker containing distilled water. The finger was then removed to allow the test flour sample to be dumped into distilled water. The wettability was recorded as the time required for the sample to be completely wet.

### **VISCOSITY**

The rotating spindle method described by [13] was employed in the viscosity determination. The

viscosities of each of the sample were determined with a viscometer. 5 g of the sample was dissolved in 100 mL of water in a disposable plastic cup. The cup with its content was placed in a water bath and heated up to boiling. Each sample was in a separate cup. The cups were then removed and cooled to room temperature of about 25 °C. Each sample in the disposable cup was placed under the equipment at speed of 30 rpm. The viscosity was recorded in centipoises (cp).

### **STATISTICAL ANALYSIS**

All analyses were done in triplicate to evaluate experimental reproducibility and reported as Mean ± Standard Deviation. The data obtained were subjected to one way analysis of variance (ANOVA) using SPSS version 21. Duncan's multiple range test (DMRT) was used to determine means that were significantly different at a level of significance ( $\alpha$ ) of 0.05.

## **RESULTS AND DISCUSSION**

### **PROXIMATE ANALYSIS**

The result of the proximate analysis is presented in table 3.1. The moisture content of *Colocasia esculenta* flour ranges from 5.98% to 11.28% while CE90 has the lowest value (5.98%) and CE50 having the highest value (11.28%). Significant differences were observed in the obtained result. It was also observed that moisture content reduces in the flour sample as the drying temperature increases and this was in agreement with the research earlier reported by [15]. The ash content of the samples ranges from 0.75% to 3.22%. The highest value (3.22%) was recorded in CE50 and the lowest value recorded in CE90 (0.75%). Earlier research by [16] and [17] reported ash content of *Colocasia esculenta* ranges from 0.51% to 2.98% and 0.93% to 3.02% respectively, which is similar to that reported in this research. It was observed that the ash content decreases with increase in

temperature. The result indicates the presence of inorganic nutrients in the flour samples, therefore the samples could be a source of mineral elements having nutritional importance [18].

The fiber content ranges from 2.75 % to 3.19%. CE90 recorded the highest value (3.19%) while CE70 recorded the lowest value (2.75%) even though significant difference wasn't noticed in the five samples. The fat content of the cocoyam flour samples ranges from 0.93% to 0.41%. CE50 had the highest value (0.93%) while CE90 had the lowest value (0.41%). It was observed that the fat content reduces with increase in temperature as earlier reported by [19] and this may be attributed to the oxidation of fat as temperature increases [20]. The low content of fat would enhance the shelf life of the flour due to the lowered chance of rancid flavor development [21]. Hence CE90 would tend to have longer shelf life than CE50.

The crude protein content of the flour samples ranges from 7.26% to 4.93% with CE90 having the lowest (4.93%) and CE50 with the highest (7.26%). The protein content reported in this present research is slightly higher than 6.96 to 4.34% reported by [22]. It was observed that as temperature increases, the protein content decreases and this variation could be attributed to the denaturing of protein as temperature increases. The carbohydrate content of the flour samples ranges from 81.23% to 72.69%. CE90 was observed to have the highest value while CE50 had the lowest value. CE90, CE80, CE70 showed no significant difference compared to CE60 and CE50. It was observed that temperature affect the composition of carbohydrate as it increases with increase in temperature. The result reported high carbohydrate content values in *Colocasia esculenta* which is in line with the research done by [23]. Carbohydrate has been discovered to be the predominant of all the nutrients in roots and tubers [24].

**Table 3.1. Proximate Analysis of *Colocasia esculenta***

PARAMETERS						
Sample	Moisture Content (%)	Ash Content (%)	Protein	Fat	Crude fiber	Carbohydrate
CE50	11.28 <sup>a</sup> ±1.86	3.22 <sup>a</sup> ±0.91	7.26 <sup>a</sup> ±1.11	0.93 <sup>a</sup> ±0.07	2.82 <sup>b</sup> ±0.43	72.69 <sup>c</sup> ±1.71
CE60	9.82 <sup>a</sup> ±1.13	1.81 <sup>b</sup> ±0.47	6.74 <sup>a</sup> ±0.83	0.81 <sup>b</sup> ±0.04	3.08 <sup>a</sup> ±0.80	75.74 <sup>b</sup> ±1.67
CE70	8.73 <sup>b</sup> ±1.04	1.39 <sup>b</sup> ±0.55	5.51 <sup>b</sup> ±0.46	0.67 <sup>c</sup> ±0.06	2.75 <sup>b</sup> ±0.61	77.51 <sup>a</sup> ±1.95
CE80	6.51 <sup>b</sup> ±1.19	1.08 <sup>c</sup> ±0.63	5.12 <sup>b</sup> ±0.58	0.55 <sup>d</sup> ±0.02	3.13 <sup>a</sup> ±0.88	80.59 <sup>a</sup> ±2.54
CE90	5.98 <sup>c</sup> ±0.85	0.75 <sup>d</sup> ±0.27	4.93 <sup>c</sup> ±0.21	0.41 <sup>e</sup> ±0.05	3.19 <sup>a</sup> ±0.74	81.23 <sup>a</sup> ±2.31

Number of replicates = 3; Mean ± Standard Deviation; Mean with different superscript across rows are significantly different at (P<0.05)

Sample codes: *Colocasia esculenta* dried at 50 °C denoted as CE50

*Colocasia esculenta* dried at 60 °C denoted as CE60

*Colocasia esculenta* dried at 70 °C denoted as CE70

*Colocasia esculenta* dried at 80 °C denoted as CE80

*Colocasia esculenta* dried at 90 °C denoted as CE90

**FUNCTIONAL PROPERTIES**

**WATER ABSORPTION CAPACITY (WAC)**

The water absorption capacity (WAC) of the flour samples ranges from 1.32 mL/g to 2.9mL/g (Table 3.2) with CE90 having the highest and CE50 had

the lowest. There was a significant difference in the water absorption capacity of the flour samples. CE90 showed higher absorption capacity than the other samples. [25] reported a slightly close WAC values with the obtained result in this research. However, [26] recorded a higher value compared to the values recorded in this research.

It was observed that WAC increased with increase in temperature. This same observation was reported by [27] and this can be attributed to the fact that the hydrophilic tendency of the starch increased with increase in its drying temperature. Also at higher temperature, starch expands rapidly especially in the amorphous region [28]. WAC is important in the development of ready to eat foods and a high absorption capacity may assure product cohesiveness [29].

### **OIL ABSORPTION CAPACITY (OAC)**

The oil absorption capacity (OAC) of the flour samples ranges from 0.63 mL/g to 1.43 mL/g. CE90 had the highest oil absorption capacity value (1.43 mL/g) and CE50 had the lowest value (0.63 mL/g). It was observed that OAC increased with increase in temperature which could be attributed to the fact that higher temperature yielded low moisture content thereby allowing oil to be absorbed. The absorption of oil by food products improves flavour retention and mouth feel thereby giving soft texture and good flavour to food [30].

### **BULK DENSITY**

The bulk density ranges from 0.59 g/mL to 0.95 g/mL with CE50 having the highest value and CE90 had the lowest value. There was no significant difference between CE60 and CE70. The bulk density reduces with increase in temperature and this is as a result of the sample dried at high temperature had lower moisture content thereby having less surface area compare to the samples with higher moisture content. High bulk density is desirable for greater ease of dispersibility and reduction of paste thickness as it gives an indication of the relative volume of packaging material required [31].

### **SWELLING CAPACITY**

The swelling index of the flour samples varied from 14.30 to 18.60. Significant difference was

observed among CE70, CE80, and CE90 while there was no significant difference between CE50 and CE60. The values obtained in this research were more than value of cocoyam reported by [32]. It was observed that swelling capacity increases with increase in temperature and as expected also varies directly as water absorption capacity. Swelling capacity indicate strength and character of the starch granules. Generally cocoyam samples show good swelling index when compared to other root crops like cassava. This is because cocoyam has starch granules with highly digestible nature. The starch grain of cocoyam is about one tenth of potato starch grain [33].

### **WETTABILITY CAPACITY**

The flour samples have wettability capacity ranged from 12secs to 23secs with CE90 and CE80 having the lowest value (12 secs) while CE50 has the highest value (23 secs). There were significant differences in the wettability capacity of CE50, CE60, and CE70 while CE90 and CE80 did not differ significantly. It was observed that wettability reduces with increase in temperature and this could be attributed to the high temperature treatment which made them to absorb moisture faster thereby making them to have a low wetting time [34][35].

### **VISCOCITY**

The viscosity of the flour samples varied from 21 cp to 35 cp. CE90 had the highest value while CE50 had the lowest value. The viscosity of CE90 (34 cp) differed significantly from CE80 (29 cp), CE60 (26 cp) and CE50 (21cp). It was observed that viscosity increases with increase in temperature. This result implies that more viscous cocoyam flour could be obtained with further raising of the drying temperature [36]

### **GELATINIZATION POINT**

Gelatinization in food refers to the disruption of starch in which starch granules swell when heated in the presence of water [37]. The

gelatinization point of the flour samples varied from 81 °C to 90 °C with CE50 having the lowest while CE90 had the highest value. There was no significant difference in the gelling point of CE90, CE80 and CE70 although CE50 differs significantly

from CE60. The low gelatinization point of CE50 might be attributed to the low water absorption capacity of sample CE50 compared to CE90 which has a higher gelatinization point and water absorption capacity.

**Table 3.2. Functional Properties of *Colocasia esculenta***

PARAMETERS							
Sampe	WAC(mL/g)	OAC(mL/g)	BD (g/mL)	GP (°C)	SC	WC(Secs)	VM (cp)
CE50	1.32 <sup>c</sup> ±0.17	0.63 <sup>d</sup> ±0.09	0.95 <sup>a</sup> ±0.12	81 <sup>c</sup> ±2.70	14.30 <sup>d</sup> ±0.96	23 <sup>a</sup> ±3.00	21 <sup>e</sup> ±1.50
CE60	1.69 <sup>b</sup> ±0.11	0.91 <sup>c</sup> ±0.18	0.91 <sup>a</sup> ±0.09	82 <sup>b</sup> ±2.40	14.80 <sup>d</sup> ±1.11	18 <sup>b</sup> ±1.50	26 <sup>d</sup> ±1.50
CE70	1.93 <sup>b</sup> ±0.28	1.08 <sup>c</sup> ±0.05	0.87 <sup>ab</sup> ±0.09	85 <sup>ab</sup> ±1.75	15.90 <sup>c</sup> ±1.06	15 <sup>c</sup> ±2.00	32 <sup>ab</sup> ±2.50
CE80	2.44 <sup>a</sup> ±0.45	1.12 <sup>b</sup> ±0.18	0.71 <sup>b</sup> ±0.10	87 <sup>a</sup> ±2.85	16.20 <sup>b</sup> ±1.28	12 <sup>d</sup> ±1.50	29 <sup>c</sup> ±1.86
CE90	2.90 <sup>a</sup> ±0.61	1.43 <sup>a</sup> ±0.12	0.59 <sup>c</sup> ±0.08	90 <sup>a</sup> ±4.50	18.60 <sup>a</sup> ±1.56	12 <sup>d</sup> ±1.00	35 <sup>a</sup> ±3.50

Number of replicates = 3; Mean ± Standard Deviation; Mean with different superscript across rows are significantly different at (P<0.05). WAC- water absorption capacity. OAC- oil absorption capacity. BD- bulk density. GP- gelatinization point. SC- swelling capacity. WC- wettability capacity. VM- viscosity measurement

### CONCLUSION

The results of this research indicate that drying temperature have effect on the proximate and functional composition of (*Colocasia esculenta*) cocoyam flour. Depending on the storage method and intended end use of the flour, different drying temperatures can be adopted. The use of high drying temperature can be a good advantage for sample's easy transportation and storability, since high drying temperature showed low bulk density.

### ACKNOWLEDGEMENT

Dr. (Mrs) Adaramoye and Miss Kolawole Oreoluwa, thank you so much for the contribution and encouragement toward the completion of this work. Your effort can never be forgotten.

### REFERENCES

[1]. Shiyam, J. O; Obiefuna, J. C; Ofoh, M. C. Oko, B. F. D. and Uko, A. E. (2007). Growth and corm yield response of upland cocoyam (*xanthosomasagittifolium*) to sawdust mulch and n p k 20 : 10: 10 fertilizer rates in the humid forest zone of

Nigeria Continental J. Agronomy 1:5 -10, 2007, pp5-11.  
 [2]. Ekpo, M.O. (2001). Gender Implication for Sustainable Technology adaptation In: Akoroda, M.O. and Ngeve, Jim (eds) Proceedings of the 7th Triennial Symposium of the International society for Tropical Root and Tuber Crops (ISTRC). Cotonou Benin. Oct. 11- 17th, 1998: 110-120.  
 [3]. Sagoe, R.Marfo, K.A and Dankyl, A.A (2001). The potentials of cocoyam production in Ghana In: Akoroda, M.O, Ngeve and J.M(eds). Proceedings of the 7th Triennial symposium of the international society for Tropical Root and Tuber Crops (ISTRC).  
 [4]. Famurewa, J. A. V. Olatujoye, J. B. and Ajibode, A. (2014) Drying Phenomenon and Influence on the Anti Nutritional and Pasting Properties of Cocoayam (Taro). *Journal of Scientific Research & Reports*, 3(2): 275-283, Article no. JSRR.2014.002.  
 [5]. Bamgboye, A.I. and Adejumo, O.I. (2010). Thermal properties of roselle seeds, International Agrophysics vol: 24, number: 1, pages: 85-87.



- [6]. Ajewole, A.I. (2004). Enrichment of Pregelatinized Cocoyam Flour with melon seeds protein Concentrate as an Instant Food Gel. *JAT* 5(2): 28-32.
- [7]. Ayele, T. and W.K. Nip, (1994). Functional properties of raw and precooked taro (*Colocasiaesculenta*) flours. *Int. J. Food Sci. Technol.*, 29: 457-462.
- [8]. Association of Official Analytical Chemists (AOAC), (2000). Official Methods of Analysis of the Association of the Official Analytical Chemists. 18th Edn., Washington DC.
- [9]. American Association of Cereal Chemists (AACC), (2002), Approved method of the American Association of Cereal Chemists, 10th Ed., St Paul, MN, 231-249.
- [10]. Onwuka, G. I. (2005). Food Analysis and Instrumentation: Theory and Practice. Naphthali prints, Lagos. 102-106.
- [11]. Niba, L.L., Bokanga, M., Jackson, F.I., Schlimme, D.S. and Li, B.W.(2001). Physio chemical. properties and starch granular characteristics of flour from various *Manihotesculenta* (cassava) genotypes. *J. Ed.Sc.* 67: 1701.
- [12]. Abbey, B. W. and Ibeh, G.O. (1988). Functional properties of raw and heat processed cowpea flour. *J. Food Science* 53(6). 1774-1777, 1791.
- [13]. Ogunlakin, G.O. Oke, M.O., Babarinde, G.O. and Olatunbosun, D.G. (2012). Effect of Drying Methods on Proximate Composition and Physico-chemical Properties of Cocoyam Flour. *American Journal of Food Technology*, 7: 245-250.
- [14]. Ojinaka, M.C., Akobundu, E.N.T. and Iwe, M.O. (2009). Cocoyam starch modification effects on functional, sensory and cookies qualities. *Pak. J. Nutr.*, 8: 558-567.
- [15]. Amandikwa, C. (2012). Proximate and functional properties of open air, solar and oven dried cocoyam flour. *Int'l Journal of Agric. and Rural Dev.* Volume 15 (2):988-994.
- [16]. Sefa-Dede, S. and Agyir-Sackey, E.K. (2004). Chemical composition and the effect of processing on oxalate content of cocoyam *Xanthomasagittifolium* and *Colocasia esculenta*.
- [17]. Udensi, A. and Eke, O.(2000). Proximate composition and functional properties of flour produced from *Mucunacochinensis* and *Mucunautles*. In Proceedings of the 1st Annual Conference of the College of Agriculture and Veterinary Medicine Abia State University. 10-13th Sept. 2000.Pp 170-174.
- [18]. Igbabul, B. D., Amove, J., and Twadue, I. (2014). Effect of fermentation on the proximate composition, antinutritional factors and functional properties of cocoyam (*Colocasiaesculenta*) flour. *African Journal of Food Science*, 5(3), 67-74.
- [19]. Enonfom, J.A and I.B Umoh (2004). Effects of heat and tetracycline treatment on the food quality and acidity factors in cocoyam [*Xanthosoma Sagittifolium* (L) Schott]. *Park. J. Nutr* 3: 240- 243.
- [20]. Onwueme, I.C. (1982), *The Tropical Tuber Crops*. English Language Book Society, John Wiley & Sons, Chichester, UK, 1982.
- [21]. SefaDedeh and S.E Kofi-Agyir, (2002).starch structure and some properties of cocoyam [*Xanthomasagittifolium* and *Colocasiaesculenta*] starch and raphides j. food chem.; 74; 435- 444.
- [22]. Ajala A.S., Oguniola, A.D. and Odudele F. B. (2014). Evaluation of Drying Temperature on Proximate, Thermal and Physical Properties of Cocoyam Flour. *Global Journal of Engineering, design and technology.* Vol.3 (4):13-18.
- [23]. Enwere, N.J, (1998). *Foods of plant origin*. Afro-orbis publication Ltd, PP: 194- 199.
- [24]. Gbadamosi, S.O. and Oladeji, B.S (2013). Comparative studies of the functional and physico-chemical properties of isolated Cassava, Cocoyam and Breadfruit starches,

- International Food Research Journal 20(5): 2273-2277
- [25]. Asumugha, V. U. and Uwalaka, B. C. (2000). Chemical and organoleptic evaluation of snacks developed from cocoyam (*Colocasia esculenta*, *Xanthosoma affinale*) and wheat (*Triticum spp*) composite flours. *Nigerian Agricultural Journal* 31:78-88.
- [26]. Ammar, M.S, Hegazy, A.Z and Beder, S.H. (2009). Using taro flour as partial substitute of Wheat flour in bread making, *World Journal of Dairy & Food Science*, 4(2): 9-99.
- [27]. Aviara, N. A., Igbeka, J. C. and Nwokocha, L. M. (2010). Physicochemical Properties of Sorghum (*Sorghum Bicolor* L. Moench) Starch as Affected by Drying Temperature. *Agricultural Engineering International: the CIGR Ejournal*. Manuscript Vol. XII. , pp 1-5
- [28]. Adebawale, K. O; Olu-Owolabi, B. I; Olayinka, O. O. and Olayide, S. L. (2005). Effect of heat moisture treatment and annealing on physicochemical properties of red sorghum starch. *African Journal of Biotechnology*, 4(9): 923-933.
- [29]. Ihekoroonye, A. (1998). *Manual on small scale food processing*. 1st edn., Academic Publishers, Nsukka, pp: 32.
- [30]. Mbofung, C., Aboubakar, M.F., Njintang, Y.N., Bouba, A.A and Balaam, F.F. (2006). Physicochemical and functional properties of six varieties of taro (*Colocasia esculenta* L. Schott) flour, *Journal of Food Technology*, 4: 135-142.
- [31]. Oke, M.O and Bolarinwa, I.F (2012). Effect of Fermentation on Physicochemical Properties and Oxalate Content of Cocoyam (*Colocasia esculenta*) Flour. *ISRN Agronomy*, Volume 1, Article ID 978709
- [32]. Okpala, L. C. and Chinyelu, V. A. (2011). Physicochemical, Nutritional and Organoleptic evaluation of cookies from pigeon pea (*Cajanus cajan*) and cocoyam (*Xanthosoma*) flour blends. *African Journal of Food, Agriculture, Nutrition and Development* 11(6):5431-5443.
- [33]. Iwuoha, C. I., and Kalu, F. A. (1995). Calcium oxalate and physico-chemical properties of cocoyam (*Colocasia esculenta* and *Xanthosoma sagittifolium*) tuber flours as affected by processing. *Food Chemistry*, 54(1), 61-66. [http://dx.doi.org/10.1016/0308-8146\(95\)92663-5](http://dx.doi.org/10.1016/0308-8146(95)92663-5).
- [34]. Olajide, R., Akinsoyinu, A. O., Babayemi, O. J., Omojola, A. B., Abu, A. O., and Afolabi, K. D. (2011). Effect of processing on energy values, nutrient and anti-nutrient components of wild cocoyam (*Colocasia esculenta* L. Schott) corm. *Pakistan Journal of Nutrition*, 10(1), 29-34. <http://dx.doi.org/10.3923/pjn.2011.29.34>.
- [35]. Owusu-Darko, P. G., Paterson, A., and Omenyo, E. L. (2014). Cocoyam (corms and cormels) an underexploited food and feed resource. *Journal of Agricultural Chemistry and Environment*, 03(01), 22-29. <http://dx.doi.org/10.4236/jacen.2014.31004>.
- [36]. Ibarz, A., and Barbosa-Canovas, G. V. (2014). Physical and chemical properties of food powders. In A. Ibarz, & G. V. Barbosa-Canovas (Eds.), *Introduction to Food Process Engineering* (p. 261-287). Boca Raton: CRC Press. <http://dx.doi.org/10.1201/b14969-16>.
- [37]. James, E.O., Peter, I.A., Charles, N.I and Joel, N. (2013). Chemical composition and effect of processing and flour particle size on Cocoyam (*Colocasia esculenta* var. *esculenta*) flour, *Nigerian Food Journal*, 31(2): 113-122.